



FormthoticsTM
Foot Science

Is Foot Pronation a Bad Thing?

Rethinking an Old Assumption



"Pronation is not a dirty word."

Dr. John Weed



Introduction

Dr John Weed – a renowned podiatric clinician and educator – knew that pronation was often misunderstood. Yet over the years, it's become one of the most routinely "treated" foot motions in clinical and commercial settings. Orthoses, motion-control shoes, and bracing are widely prescribed with the goal of reducing or eliminating pronation.

But is pronation really pathological?
And if not always,
when is it appropriate
to intervene?



What Is Pronation?

Pronation is the motion created as the talus rotates medially and downward over the calcaneus, causing the heel to roll inward (eversion) and the medial arch to drop. This is visible as a flattened arch and valgus heel – the classic signs of “overpronation”.

This motion occurs primarily at the subtalar joint and is a natural part of walking and

running. Most individuals strike the ground in slight supination, then pronate as the foot contacts the ground fully – helping the body absorb shock and adapt to terrain.

Without pronation, the limb would receive excessive impact. In fact, high-arched feet that do not pronate well are more commonly associated with overuse injuries.



A photograph of a person's legs and feet against a light blue background. The person is sitting, and their right foot is being massaged by two hands. The left foot is flat on the ground. The skin is dark brown.

Why Does the Foot Pronate?

Pronation plays a functional role in gait. It enables the foot to adapt to uneven surfaces, absorb impact, and stabilise the body during stance. Several factors influence the degree and speed of pronation:

- Body weight and joint alignment (e.g. genu valgum)
- Tibial or femoral torsion
- Tight calf musculature limiting dorsiflexion
- Forefoot varus or instability
- Hypermobility (e.g. Ehlers-Danlos Syndrome)
- Age-related laxity or degenerative changes

The subtalar joint responds to both intrinsic and extrinsic forces. A person may pronate excessively as compensation for a forefoot deformity or in response to limb rotation, instability, or reduced range of motion elsewhere.

When Is Pronation a Problem?

While pronation is a natural motion, excessive or uncontrolled pronation can contribute to tissue stress and overuse injury – particularly when it exceeds the tissue's capacity to adapt. Conditions frequently associated with abnormal pronation include:

- Plantar fasciitis
- Medial tibial stress syndrome (MTSS)
- Achilles tendinopathy (less direct association)
- Tibial and navicular stress fractures
- Patellofemoral pain
- Iliotibial band syndrome (ITBS)
- Hallux limitus, bunions, metatarsalgia

The challenge is determining when pronation is contributing to symptoms. This depends on joint structure, gait mechanics, activity level, and individual tissue resilience.



How Orthoses Help

Foot orthoses are designed not to eliminate pronation, but to manage its degree, velocity, and impact on other joints.

They work by:

- Redistributing plantar load
- Reducing excessive midfoot collapse
- Stabilising the subtalar joint and medial arch
- Controlling tibial rotation that affects the knee

Formthotics offer customisable support that can be adjusted based on forefoot structure, arch compliance, calf length, and footwear profile – allowing clinicians to target interventions more effectively.





Children and Flat Feet

Flat feet in young children are often benign. In infancy and early childhood, a thick plantar fat pad and ligamentous laxity make the arch appear absent. Most children develop a visible arch by age 10 without intervention.

Still, orthoses may be appropriate when:

- The child is symptomatic (e.g., pain, fatigue)
- There is developmental delay or hypotonia
- Structural hypermobility impairs gait

In these cases, orthoses may help improve comfort, stability, and participation in physical activity.



Muscle Strength and Orthotic Use

A common concern is that orthoses might weaken the feet. However, EMG studies show that intrinsic and extrinsic foot muscles remain active – and may even become more efficient – with orthotic use.

Much like glasses don't weaken your eyes, orthoses don't inherently weaken your feet. Instead, they provide external support to modulate mechanical stress while allowing for ongoing functional activity.

Foot Pronation and the Knee

Because the talus transmits movement directly to the tibia, excessive pronation introduces internal tibial rotation. This movement, when paired with a knee joint designed primarily for flexion and extension, can create rotational stress – especially on the patellofemoral joint.

Foot orthoses have been shown to reduce tibial rotation and are effective in managing peri-patellar symptoms and medial knee arthritis in certain populations.



Conclusion

Pronation is not inherently problematic. It's a vital, shock-absorbing component of human locomotion. However, when excessive or poorly controlled, it may lead to overload injuries in the foot, leg, and knee.

The clinical question is not whether pronation is "bad," but whether its presentation in a given patient is contributing to symptoms – and whether targeted orthotic intervention can help.

In that respect, orthoses – particularly adaptable options like Formthotics – remain a valuable, evidence-informed tool in the conservative management of pronation-related issues. As clinicians, we can honour Dr. John Weed's legacy by not pathologising pronation, but instead understanding when and how to support it.



References

1. **Thomas et al.** The population prevalence of foot and ankle pain in middle and old age: a systematic review. *Pain*. 2011;152
2. **Mickle et al.** Foot pain, plantar pressures, and falls in older people: a prospective study. *Journal of the American Geriatric Society*. 2010;58
3. **Menz et al.** Foot Pain and Mobility Limitations in Older Adults: The Framingham Foot Study. *Journal of Gerontology*. October, 2013
4. **Davis, et al.** Effect of Inverted Orthoses on Lower-Extremity Mechanics in Runners. *Medicine and Science in Sport and Exercise*. December, 2003
5. **Cornwall & McPoil.** Footwear and Foot Orthotic Effectiveness Research: A New Approach. *Foot/Ankle Therapy and Research*. June, 1995
6. **Rabusin CL, et al.** Efficacy of heel lifts versus calf muscle eccentric exercise for mid-portion achilles tendinopathy: a randomised trial. *British Journal of Sports Medicine*, May 2021
7. **Lee KKW, Ling SKK, Yung PSH.** Controlled trial to compare the achilles tendon load during running in flatfoot participants using a customized arch support orthoses vs an orthotic heel lift. *BMC Musculoskeletal Disorders* Nov. 2019
8. **Bonanno DR, et al.** Risk Factors for Lower Limb Injuries during Naval Training: a Retrospective study. *Journal of the Royal Army Medical Corps*. Sept 2018
9. **Naderi, A, et al.** Foot orthoses enhance the effectiveness of exercise, shockwave, and ice therapy in the management of MTSS. *Clinical Journal of Sports Medicine*. March 2022
10. **Yeung, S. et al.** Interventions for Preventing Lower Limb Soft-Tissue Running Injuries. *Cochrane Library*. July 2011
11. **Kayll, SA et al.** Do biomechanical foot-based interventions reduce patellofemoral joint loads in adults with and without patellofemoral pain or osteoarthritis? A systematic review and meta-analysis. *British Journal of Sports Medicine*. July 2023
12. **Boldt et al.** effects of medially wedge foot orthoses on knee and hip joint kinematics in females with and without patellofemoral pain syndrome. *Journal of Applied Biomechanics*. February 2013
13. **McPoil et al.** Effect of Foot Orthoses Contour on Pain Perception in Individuals with Patellofemoral Pain. *Journal of the American Podiatric Medical Association*. January 2011
14. **Tan et al.** Associations of foot and ankle characteristics with knee symptoms and function in individuals with patellofemoral osteoarthritis. *Journal of Foot and Ankle Research*. September 2020
15. **Mullaji A, Shetty GM.** Persistent hindfoot valgus causes lateral deviation of weightbearing axis after total knee arthroplasty. *Clinical Orthopaedics and Related Research*. 2011;469(4)
16. **Murley, et al.** Do Foot Orthoses Change Lower Limb Muscle Activity in Low Arched Feet Towards a Pattern Observed in Normal Arched Feet? *Clinical Biomechanics*. Volume 25, Issue 7. August 2010
17. **Christensen, BH et al.** Foot Orthotics Reduce The Navicular Drop - A Novel Method Allowing For In-Shoe Measurement. *Aalborg University Hospital*. February 2014.
18. **Kilmartin, T.** The Orthotic Treatment of Juvenile Hallux Valgus. Submitted as a Doctoral Thesis, *Univeristy of Nottingham*. 1994
19. **Paquette, M.** Moving Beyond Weekly Distance: Optimizing Quantification of Training Load in Runners. *Journal of Orthopaedic & Sports Physical Therapy*. Vol 50. October, 2020
20. **Reeves, J. et al.** A Systematic Review of the Effect of Footwear, Foot Orthoses and Taping on Lower Limb Muscle Activity During Walking and Running. *Prosthetics and Orthotics International*. Vol 43, 2019
21. **Robb, K, Howe, E, Perry, S.** The effects of foot orthoses and sensory facilitation on lower limb electromyography: A scoping Review. *The Foot*. Vol 52, 2022



26 Dakota Crescent, Wigram,
Christchurch, 8042, New Zealand

www.formthotics.com